



An EnviroMax® Solution for Difficult Turf Problems



EnviroMax® IndoxaGuard 200SC Insecticide (222 g/L Indoxacarb (S:R 9:1))

A Safer Effective Alternative



COMPETITIVE BENEFITS

- EPA Designated “Reduced Risk” Insecticide.
- New and unique Mode of Action Non-repellent.
- No Cross-Resistance with other insecticide classes.
- Effective Resistance Management.
- Contact and Gut Action.
- Translaminar Activity – penetrates plant tissue providing control of pests on all surfaces.
- Broad-spectrum control. Effective against a wide range of Lepidopteran pests.
- Quality Assured - Superior Suspension Concentrate Formulation.
- Improved Application of Indoxacarb Active.
- Uniform Micron-sized Indoxacarb particles.
- Increased Insect Contact and Efficacy.
- Quality Assured Superior Suspension Concentrate Formulation.
- Significantly Reduced Sedimentation and Blocked Nozzles.
- Consistent Quality is Our Guarantee.
- Large pack size range (250mL - 20L).

High Quality HDPE Packs

- International Standard in Turf Management Formulations.
- Produced by Professionals for Professionals!
- Give Your Business the EnviroMax® Advantage!
- Competitive Pricing!
- Quality Assured!

Protect Your Turf and Ornamentals with The Best

Directions for Use and Application Situations

Situation	Pest	Rate	Critical Comments
Golf greens and surrounds.	Argentine Stem Weevil larvae and adults. (<i>Listronotus bonariensis</i>).	1.25 L/ha.	Two seasonal treatment opportunities should be targeted. Apply late winter/early spring when overwintered adults are observed to minimise egg laying and population build up. Early application is essential. Make a second application early summer as a treatment for adults and foliar feeding larvae.
Golf greens and surrounds, Fairways.	Caterpillars including Black Cutworm. (<i>Agrotis ipsilon</i>), Lawn Armyworm (<i>Spodoptera mauritia</i>).	375 mL/ha.	Apply immediately when pest activity is noted. To ensure optimum control, delay watering (irrigation) or mowing for 24 hours after application.

Mandatory downwind buffer zones (metres).

Application rate.	Boom height above the target canopy	Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to maximum label rate (1.25 L/ha).	0.5 m or lower.	0	0	0	0	0
	1.0 m or lower.	0	20	10	0	40
375 mL/ha or lower.	0.5 m or lower.	0	0	0	0	0
	1.0 m or lower.	0	5	0	0	10

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

PROTECTION OF HONEYBEES AND OTHER INSECT POLLINATORS

Highly toxic to bees. Harmful to bee brood. DO NOT allow spray drift to flowering weeds or flowering crops in the vicinity of the treatment area. Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar and pollen, if there is potential for managed hives to be affected by the spray or spray drift.

PROTECTION OF NON-TARGET BENEFICIAL INSECTS

Beneficial insects contribute to control of secondary pest outbreaks. EnviroMax® Indoxaguard applications are unlikely to affect spiders and lacewings. Applications MAY temporarily reduce populations of predatory beetles, transverse ladybirds, ants and pirate bugs, but populations quickly recover.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple rinse containers before disposal.

Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage the eyes. May irritate the skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container, mixing and loading, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and inside mouth. IF IN EYES: Wash immediately with water. Remove contact lenses, if present and easy to do so. Continue rinsing. If eye irritation persists: Get medical attention.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

In a transport emergency dial 000, Police or Fire Brigade.

APVMA Approval No.: 94130

Note: This Technical Data Sheet is for reference only. When using EnviroMax® IndoxaGuard 200SC Insecticide refer to the registered label.

Active Constituent: 222 g/L Indoxacarb S:R 9:1.

Inert ingredients: Other nonhazardous ingredients up to 100%.

Formulation Type: Aqueous Suspension Concentrate (SC).

Pesticide Group: Oxadiazine.

Insecticide Mode of Action: Sodium Channel Blocker.

EnviroMax® Technologies Pty Ltd

A PO Box 305, North Lakes, Queensland 4509, Australia.

T +61 7 3088 9111 E info@enviromaxtechnologies.com.au

www.enviromaxtechnologies.com.au

EnviroMax® is a registered trademark of EnviroMax® Technologies Pty Ltd, Australia.

